

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

35-1174: Polyclonal Antibody to Histone H3.1 (Phospho-Ser10)

Clonality: Polyclonal Application: WB,IHC,IF

Reactivity: Human, Mouse, Rat

 Gene :
 HIST1H3A

 Gene ID :
 8350

 Uniprot ID :
 P68431

 Format :
 Purified

 Alternative Name :
 H3/b,H3FB

 Isotype :
 Rabbit IgG

Immunogen Information : Peptide sequence around phosphorylation site of serine 10 (R-K-S(p)-T-G) derived from Human

Histone H3.1.

Description

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. Dai J, et al. (2005) Genes Dev 19(4): 472-488. Yih LH, et al. (2005) Carcinogenesis 26(1): 53-63.

Product Info

Amount : 50 μl / 100 μl

Content: Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM

NaCl, 0.02% sodium azide and 50% glycerol.

Storage condition:

Storage condition:

Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Predicted MW: 17kd, Western blotting: 1:500~1:1000, Immunohistochemistry: 1:50~1:100, Immunofluorescence: 1:100~1:200




Figure 1: Immunofluorescence staining of methanol-fixed Hela cells using Histone H3.1 (Phospho-Ser10) Antibody 35-1174.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using Histone H3.1(Phospho-Ser10) Antibody 35-1174 (left) or the same antibody preincubated with blocking peptide(right).




Figure 3: Western blot analysis of extracts from SK-BR-3 cells, treated with insulin and EGF, and pretreated with U0126 and LY294002 cells using Histone H3.1 (Phospho-Ser10) Antibody 35-1174.




Figure 4: Immunofluorescence staining of methanol-fixed Hela cells using Histone H3.1 (Phospho-Ser10) Antibody 35-1174.