

35-1201: Polyclonal Antibody to FAK (Phospho-Tyr397)

Clonality :	Polyclonal
Application :	WB
Reactivity :	Human, Mouse, Rat
Gene :	PTK2
Gene ID :	5747
Uniprot ID :	Q05397
Format :	Purified
Alternative Name :	FAK, FAK1, PTK2
Isotype :	Rabbit IgG
Immunogen Information :	Peptide sequence around phosphorylation site of tyrosine 397(D-D-Y(p)-A-E) derived from Human FAK.

Description

Non-receptor protein-tyrosine kinase implicated in signaling pathways involved in cell motility, proliferation and apoptosis. Activated by tyrosine-phosphorylation in response to either integrin clustering induced by cell adhesion or antibody cross-linking, or via G-protein coupled receptor (GPCR) occupancy by ligands such as bombesin or lysophosphatidic acid, or via LDL receptor occupancy. Microtubule-induced dephosphorylation at Tyr-397 is crucial for the induction of focal adhesion disassembly. Plays a potential role in oncogenic transformations resulting in increased kinase activity.

Product Info

Amount :	50 µl / 100 µl
Content :	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage condition :	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note

Predicted MW: 125kd, Western blotting: 1:500~1:1000




Figure 1: Western blot analysis of extract from 293 cells untreated or treated with EGF using FAK(Phospho-Tyr397) Antibody using 35-1201