

## 35-1202: Polyclonal Antibody to Pyk2 (Phospho-Tyr402)

<b>Clonality :</b>	Polyclonal
<b>Application :</b>	WB,IHC,IF
<b>Reactivity :</b>	Human,Mouse,Rat
<b>Gene :</b>	PTK2B
<b>Gene ID :</b>	2185
<b>Uniprot ID :</b>	Q14289
<b>Format :</b>	Purified
<b>Alternative Name :</b>	FADK 2, FAK2, Focal adhesion kinase 2, PTK2B, Proline-rich tyrosine kinase 2 RAFTK
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	Peptide sequence around phosphorylation site of tyrosine 402 (D-I-Y(p)-A-E) derived from Human Pyk2.

### Description

Involved in calcium induced regulation of ion channel and activation of the map kinase signaling pathway. May represent an important signaling intermediate between neuropeptide activated receptors or neurotransmitters that increase calcium flux and the downstream signals that regulate neuronal activity. Interacts with the SH2 domain of Grb2. May phosphorylate the voltage-gated potassium channel protein Kv1.2. Its activation is highly correlated with the stimulation of c-Jun N-terminal kinase activity. Involved in osmotic stress-dependent SNCA 'Tyr-125' phosphorylation.

### Product Info

<b>Amount :</b>	50 µl / 100 µl
<b>Content :</b>	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
<b>Storage condition :</b>	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

### Application Note

Predicted MW: 116kd, Western blotting: 1:500~1:1000, Immunohistochemistry: 1:50~1:100, Immunofluorescence: 1:100~1:200




Figure 1: Western blot analysis of extracts from 293 cells untreated or treated with Serum using Pyk2(Phospho-Tyr402) Antibody 35-1202 .




Figure 2: Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using Pyk2(Phospho-Tyr402) Antibody 35-1202 (left) or the same antibody preincubated with blocking peptide(right).




Figure 3: Immunofluorescence staining of methanol-fixed Hela cells using Pyk2(Phospho-Tyr402) Antibody 35-1202 .