

### 35-1275: Polyclonal Antibody to IKK-beta(Phospho-Tyr199)

<b>Clonality :</b>	Polyclonal
<b>Application :</b>	IF
<b>Reactivity :</b>	Rat,Mouse,Human
<b>Gene :</b>	IKBKB
<b>Gene ID :</b>	3551
<b>Uniprot ID :</b>	O14920
<b>Format :</b>	Purified
<b>Alternative Name :</b>	IKK2, IKKB, NFKB1KB, IKK-beta
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	Peptide sequence around phosphorylation site of tyrosine 199(Q-K-Y(p)-T-V)derived from Human IKK-B.

#### Description

Serine kinase that plays an essential role in the NF-kappa-B signaling pathway which is activated by multiple stimuli such as inflammatory cytokines, bacterial or viral products, DNA damages or other cellular stresses. Acts as part of the canonical IKK complex in the conventional pathway of NF-kappa-B activation and phosphorylates inhibitors of NF-kappa-B on 2 critical serine residues. These modifications allow polyubiquitination of the inhibitors and subsequent degradation by the proteasome. In turn, free NF-kappa-B is translocated into the nucleus and activates the transcription of hundreds of genes involved in immune response, growth control, or protection against apoptosis. In addition to the NF-kappa-B inhibitors, phosphorylates several other components of the signaling pathway including NEMO/IKBKG, NF-kappa-B subunits RELA and NFkB1, as well as IKK-related kinases TBK1 and IKBKE. IKK-related kinase phosphorylations may prevent the overproduction of inflammatory mediators since they exert a negative regulation on canonical IKKs. Also phosphorylates other substrates including NCOA3, BCL10 and IRS1. Within the nucleus, acts as an adapter protein for NFKBIA degradation in UV-induced NF-kappa-B activation.

#### Product Info

<b>Amount :</b>	50 µl / 100 µl
<b>Content :</b>	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
<b>Storage condition :</b>	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

#### Application Note

Predicted MW: 87kd, Immunofluorescence: 1:100~1:200




Figure 1: Immunofluorescence staining of methanol-fixed HeLa cells using IKK-b(Phospho-Tyr199) Antibody (35-1275 , Red).