

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

35-1313: Polyclonal Antibody to Aurora A (phospho-Thr288)

Clonality: Polyclonal
Application: IHC,WB
Reactivity: Mouse,Human

Gene: AURKA
Gene ID: 6790
Uniprot ID: 014965
Format: Purified

Alternative Name: AIK, ARK1, AURA, BTAK, STK6

Isotype: Rabbit IgG

Immunogen Information: Peptide sequence around phosphorylation site of threonine 288 (R-T-T(p)-L-M) derived from

Human Aurora A.

Description

Contributes to the regulation of cell cycle progression. Required for normal mitosis. Associates with the centrosome and the spindle microtubules during mitosis and functions in centrosome maturation, spindle assembly, maintenance of spindle bipolarity, centrosome separation and mitotic checkpoint control. Phosphorylates numerous target proteins, including ARHGEF2, BRCA1, KIF2A, NDEL1, PARD3, PLK1 and BORA. Regulates KIF2A tubulin depolymerase activity By similarity. Required for normal axon formation. Plays a role in microtubule remodeling during neurite extension. Important for microtubule formation and/or stabilization. Bischoff, J.R. et al. (1998) EMBO J 17, 3052-65. Hauf, S. et al. (2003) J Cell Biol 161, 281-94. Walter, A.O. et al. (2000) Oncogene 19, 4906-16. Zhou, H. et al. (1998) Nat Genet 20, 189-93.

Product Info

Amount : 50 μl / 100 μl

Content: Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM

NaCl, 0.02% sodium azide and 50% glycerol.

Storage condition : Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Predicted MW: 48kd, Immunohistochemistry: 1:50~1:100




Figure 1: Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using Aurora A(Phospho-Thr288) Antibody 35-1313 (left) or the same antibody preincubated with blocking peptide(right).



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Western blot analysis of extracts from HT29 cells, treated with Hydroxyurea or calf intestinal phosphatase (CIP), using Aurora A (phospho-Thr288) Antibody 35-1313 .