

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 35-1330: Polyclonal Antibody to MDM2 (phospho-Ser166)

Clonality: Polyclonal Application: WB,IHC,IF Reactivity: Human Gene: MDM<sub>2</sub> Gene ID: 4193 **Uniprot ID:** 000987 Format: Purified **Alternative Name:** HDMX,hdm2 Rabbit IgG Isotype:

Immunogen Information: Peptide sequence around phosphorylation site of Serine 166 (A-I-S(p)-E-T) derived from

Human MDM2.

## **Description**

This gene is a target gene of the transcription factor tumor protein p53. The encoded protein is a nuclear phosphoprotein that binds and inhibits transactivation by tumor protein p53, as part of an autoregulatory negative feedback loop. Overexpression of this gene can result in excessive inactivation of tumor protein p53, diminishing its tumor suppressor function. This protein has E3 ubiquitin ligase activity, which targets tumor protein p53 for proteasomal degradation. This protein also affects the cell cycle, apoptosis, and tumorigenesis through interactions with other proteins, including retinoblastoma 1 and ribosomal protein L5. More than 40 different alternatively spliced transcript variants have been isolated from both tumor and normal tissues

## **Product Info**

**Amount :** 50 μl / 100 μl

Content: Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM

NaCl, 0.02% sodium azide and 50% glycerol.

**Storage condition :** Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

## **Application Note**

Predicted MW: 90kd, Western blotting:  $1:500\sim1:1000$ , Immunohistochemistry:  $1:50\sim1:100$ , Immunofluorescence:  $1:100\sim1:200$ 




Figure 1: Western blot analysis of extracts from 293 cells untreated or treated with Hydroxyurea using MDM2(phospho-Ser166) Antibody 35-1330 .



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using MDM2(Phospho-Ser166) Antibody 35-1330 (left) or the same antibody preincubated with blocking peptide(right).




Figure 3: Immunofluorescence staining of methanol-fixed Hela cells using MDM2(phospho-Ser166) Antibody 35-1330.