

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

35-1458: Polyclonal Antibody to ICAM-1 (Ab-512)

Clonality: Polyclonal

Application: WB Reactivity: Human Gene: ICAM1 Gene ID: 3383 **Uniprot ID:** P05362 Format: Purified **Alternative Name:** ICA1, ICAM1 Isotype: Rabbit IgG

Immunogen Information: Peptide sequence around aa. 510~514 (K-K-Y-R-L) derived from Human ICAM-1.

Description

ICAM proteins are ligands for the leukocyte adhesion protein LFA-1 (integrin a-L/beta-2). During leukocyte trans-endothelial migration, ICAM1 engagement promotes the assembly of endothelial apical cups through SGEF and RHOG activation. In case of rhinovirus infection acts as a cellular receptor for the virus. Greenwood J, et al. (2003) J Immunol; 171(4):2099-2108. Zhou Z, et al. (2005) Eur J Pharmacol; 513(1-2):1-8. Chen YH, et al. (2001) J Cell Biochem; 82(3):512-521

Product Info

Amount: 50 μl / 100 μl

Content: Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM

NaCl, 0.02% sodium azide and 50% glycerol.

Storage condition : Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Predicted MW: 89 92kd, Western blotting: 1:500~1:1000




Figure 1: Western blot analysis of extracts from MDA cells using ICAM-1 (Ab-512) Antibody 35-1458.