

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

35-1489: Polyclonal Antibody to HDAC4 (Ab-632)

Clonality: Polyclonal Application: WB,IF

Reactivity: Human, Mouse, Rat

Gene : HDAC4
Gene ID : 9759
Uniprot ID : P56524
Format : Purified

Alternative Name: HD4, Histone deacetylase 4

Isotype: Rabbit IgG

Immunogen Information: Peptide sequence around aa. 630~634 (A-Q-S-S-P) derived from Human HDAC4.

Description

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Involved in muscle maturation via its interaction with the myocyte enhancer factors such as MEF2A, MEF2C and MEF2D.

Product Info

Amount : 50 μl / 100 μl

Content: Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM

NaCl, 0.02% sodium azide and 50% glycerol.

Storage condition : Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Predicted MW: 140kd, Western blotting: 1:500~1:1000, Immunofluorescence: 1:100~1:200




Figure 1: Western blot analysis of extracts from HepG2 cells using HDAC4 (Ab-632) Antibody 35-1489.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Immunofluorescence staining of methanol-fixed Hela cells using HDAC4 (Ab-632) Antibody 35-1489 .