

## 35-1492: Polyclonal Antibody to cdc25C (Ab-216)

<b>Clonality :</b>	Polyclonal
<b>Application :</b>	WB,IHC,IF
<b>Reactivity :</b>	Human
<b>Gene :</b>	CDC25C
<b>Gene ID :</b>	995
<b>Uniprot ID :</b>	P30307
<b>Format :</b>	Purified
<b>Alternative Name :</b>	CDC25M1, MIP13, Dual specificity phosphatase Cdc25C
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	Peptide sequence around aa.214~218 (S-P-S-M-P) derived from Human cdc25C.

### Description

cdc25C is highly conserved during evolution and it plays a key role in the regulation of cell division. The encoded protein is a tyrosine phosphatase and belongs to the Cdc25 phosphatase family. It directs dephosphorylation of cyclin B-bound CDC2 and triggers entry into mitosis. It is also thought to suppress p53-induced growth arrest. Multiple alternatively spliced transcript variants of this gene have been described, however, the full-length nature of many of them is not known.

### Product Info

<b>Amount :</b>	50 µl / 100 µl
<b>Content :</b>	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
<b>Storage condition :</b>	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

### Application Note

Predicted MW: 60kd, Western blotting: 1:500~1:1000, Immunohistochemistry: 1:50~1:100, Immunofluorescence: 1:100~1:200




Figure 1: Western blot analysis of extracts from HT29 cells using cdc25C (Ab-216) Antibody 35-1492 and the same antibody preincubated with blocking peptide.




Figure 2: Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using cdc25C (Ab-216) Antibody 35-1492 (left) or the same antibody preincubated with blocking peptide(right).




Figure 3: Immunofluorescence staining of methanol-fixed Hela cells using cdc25C(Ab-216) Antibody 35-1492 .