

## 35-1678: Polyclonal Antibody to OCT4

<b>Clonality :</b>	Polyclonal
<b>Application :</b>	WB
<b>Reactivity :</b>	Human
<b>Gene :</b>	POU5F1
<b>Gene ID :</b>	5460
<b>Uniprot ID :</b>	Q01860
<b>Format :</b>	Purified
<b>Alternative Name :</b>	Otc3, OTF3, OTF4, OTF-3
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	Peptide sequence around aa.232~236 (R-K-R-T-S) derived from OCT-4

### Description

Transcription factor that binds to the octamer motif (5'-ATTGTCAT-3'). Forms a trimeric complex with SOX2 on DNA and controls the expression of a number of genes involved in embryonic development such as YES1, FGF4, UTF1 and ZFP206. Critical for early embryogenesis and for embryonic stem cell pluripotency Pesce, M. and Scholer, H.R. (2001) Stem Cells 19, 271-278. Pan, G. and Thomson, J.A. (2007) Cell Res 17, 42-9. Cauffman, G. et al. (2006) Stem Cells 24, 2685-91.

### Product Info

<b>Amount :</b>	50 µl / 100 µl
<b>Content :</b>	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
<b>Storage condition :</b>	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

### Application Note

Predicted MW: 45kd, Western blotting: 1:1000




Figure 1: Western blot analysis of extracts from human Umbilical cord mesenchymal stem cell using OCT-4 Antibody 35-1678 .