

35-1740: Polyclonal Antibody to STAT2 (Ab-690)

Clonality :	Polyclonal
Application :	WB,IHC
Reactivity :	Human
Gene :	STAT2
Gene ID :	6773
Uniprot ID :	P52630
Format :	Purified
Alternative Name :	P113, ISGF-3, STAT113
lsotype :	Rabbit IgG
Immunogen Information : Peptide sequence around aa.688~692 (R-K-Y-L-K) derived from Human STAT2.	

Description

STAT2 encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. In response to interferon (IFN), this protein forms a complex with STAT1 and IFN regulatory factor family protein p48 (ISGF3G), in which this protein acts as a transactivator, but lacks the ability to bind DNA directly. Transcription adaptor P300/CBP (EP300/CREBBP) has been shown to interact specifically with this protein, which is thought to be involved in the process of blocking IFN-a response by adenovirus.

Product Info

Amount :	50 μl / 100 μl
Content :	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage condition :	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note

Predicted MW: 113kd, Western blotting: 1:500~1:1000, Immunohistochemistry: 1:50~1:100



Figure 1: Western blot analysis of extracts from Hela cells using STAT2(Ab-690) Antibody 35-1740 and the same antibody preincubated with blocking peptide.

₩ abeomics

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com



Figure 2: Immunohistochemical analysis of paraffin-embedded human lung carcinoma tissue using STAT2(Ab-690) Antibody 35-1740 (left) or the same antibody preincubated with blocking peptide(right).