

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

35-1813: Polyclonal Antibody to Gamma-Catenin Antibody

Clonality: Polyclonal

Application: WB

Reactivity: Mouse, Human

 Gene :
 JUP

 Gene ID :
 3728

 Uniprot ID :
 P14923

 Format :
 Purified

Alternative Name: DP3, PDGB, PKGB, CTNNG, ARVD12

Isotype: Rabbit IgG

Immunogen Information: Peptide sequence around aa.726~730(I-D-T-Y-S) derived from Human g-Catenin

Description

Common junctional plaque protein. The membrane-associated plaques are architectural elements in an important strategic position to influence the arrangement and function of both the cytoskeleton and the cells within the tissue. The presence of plakoglobin in both the desmosomes and in the intermediate junctions suggests that it plays a central role in the structure and function of submembranous plaques. Acts as a substrate for VE-PTP and is required by it to stimulate VE-cadherin function in endothelial cells. Can replace beta-catenin in E-cadherin/catenin adhesion complexes which are proposed to couple cadherins to the actin cytoskeleton "Characterization of the interactions of a-catenin with a-actinin and beta-catenin/plakoglobin." Nieset J.E., Redfield A.R., Jin F., et al. J. Cell Sci. 110:1013-1022(1997) "Interaction of the DF3/MUC1 breast carcinoma-associated antigen and beta-catenin in cell adhesion." Yamamoto M., Bharti A., Li Y., Kufe D. J. Biol. Chem. 272:12492-12494(1997) "The transmembrane receptor protein tyrosine phosphatase DEP1 interacts with p120(ctn)." Holsinger L.J., Ward K., Duffield B., Zachwieja J., Jallal B. Oncogene 21:7067-7076(2002)

Product Info

Amount: 50 μl / 100 μl

Content: Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM

NaCl, 0.02% sodium azide and 50% glycerol.

Storage condition:

Storage condition:

Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Predicted MW: 83kd, Western blotting: 1:500~1:1000



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 1: Western blot analysis of extracts from HT-29, MCF-7 and C2C12 cells using g-Catenin Antibody 35-1813 .