

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 34-1018: Monoclonal Antibody to Calbindin(Clone: 4H7)

Clonality: Monoclonal

Clone Name: 4H7
Application: WB
Reactivity: Human
Gene: CALB1
Gene ID: 793
Uniprot ID: P05937
Format: Purified

Alternative Name: Calbindin D28,D-28K,Vitamin D-dependent calcium-binding protein, avian-type

**Isotype:** Mouse, IgG1

Immunogen Information: Full-length recombinant human protein

## **Product Info**

**Amount :** 50 μl / 100 μl

**Content:** Antibody is supplied as an aliquot of 1 mg/ml of affinity purified antibody

**Storage condition :** Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid

repeated freeze and thaw cycles.

## **Application Note**

WB: 1:1,000-1:5,000. IF/ICC or IHC: 1:1,000.




Figure-1: Immunofluorescent analysis of rat brain cerebellum section stained with mouse mAb to calbindin,(34-1018), dilution 1:1,000, in green, and costained with rabbit pAb to calretinin, dilution 1:5,000 in red. The blue is DAPI staining of nuclear DNA Following transcardial perfusion with 4% paraformaldehyde, brain was post fixed for 24 hours, cut to 45½4M, and free-floating sections were stained with the above antibodies. Calbindin antibody prominently labels the dendrites and perikarya of Purkinje cells in the molecular layer of cerebellum. In contrast the calretinin antibody stains granule cells, in the granualar layer, and their processes in the molecular layer.




Figure-2: Western blot analysis of different neuronal tissue lysates using mouse mAb to calbindin,(34-1018), dilution 1:2,000: [1] protein standard, [2] rat cerebellum, [3] pig hippocampus, and [4] cow cerebellum. Band at  $\sim$ 25 kDa corresponds to calbindin protein, heavily expressed in the cerebellum but less in hippocampus.