

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

34-1053: Monoclonal Antibody to a-internexin/NF66 (Clone: 2E3)

Clonality: Monoclonal

Clone Name: 2E3

Application: WB, IF/ICC, IHC

Reactivity: Human, Rat, Mouse, Cow

 Gene :
 INA

 Gene ID :
 9118

 Uniprot ID :
 Q16352

 Format :
 T.C. Sup.

Alternative Name: 66 kDa neurofilament protein, Neurofilament 5

Isotype: Mouse, IgG1

Immunogen Information: Purified recombinant rat Alpha -internexin expressed in and purified from E. coli.

Product Info

Amount : 50 μl / 500 μl

Content: Antibody is supplied as an aliquot of 1 mg/ml of affinity purified antibody or tissue culture

supernatant.

Storage condition:

Storage condition:

Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

WB: 1:10,000. IF/ICC and IHC: 1:5,000.




Figure-1: Immunohistochemistry of a section of rat facial nucleus 7 days following axotomy. These neurons are capable of regenerating their axons and also, concomitant with regeneration, strongly upregulate α -internexin in their perikarya. Other central neurons which are not able to regenerate their axons do not upregulate this protein after axotomy and untreated facial neurons normally show only very low levels of α -internexin. Both findings suggest that α -internexin has a role in axonal regeneration.




Figure-2: Western blot analysis of different tissue lysates using mouse mAb to α -internexin,(34-1053), dilution 1:10,000 in red: [1] protein standard, [2] rat brain, [3] rat spinal cord, [4] mouse brain, [5] mouse spinal cord, and [6] cow spinal cord lysate. The (34-1053) antibody reveals the α -internexin protein with an apparent molecular weight of 64-66kDa, with some variability across species.