

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

10-9504-B: Biotinylated Recombinant Rabbit Monoclonal Antibody to Mouse IgG1 (Clone: RM106)(Discontinued)

Clonality: Monoclonal Clone Name: RM106

Application : IP,ELISA,IHC,FACS,WB

Reactivity: Mouse Conjugate: Biotin Gene: lghg1 Gene ID: 16017 **Uniprot ID:** P01868 Format: Purified **Alternative Name:** Ighg1, Igh-4 Isotype: Rabbit IgG Immunogen Information: Mouse IgG

Product Info

Amount: 50 μg

Purification: Protein A affinity purified from an animal origin-free culture supernatant **Content:** 1 mg/ml in 50% Glycerol/PBS with 1% BSA and 0.09% sodium azide

Storage condition: Store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note

Clone RM106 reacts to the Fc region of mouse lgG1. No cross reactivity with mouse lgG2a, lgG3, lgM, lgA, lgE, human lgG, or rat lgG. The Fc region of Clone RM106 has been engineered to eliminate Fc receptor binding. ELISA: 0.005 μ g/ml-0.2 μ g/ml; Immunocytochemistry (ICC): 0.5 μ g/ml-2 μ g/ml; Immunohistochemistry (IHC): 0.5 μ g/ml-2 μ g/ml; Western Blot (WB): 0.1 μ g/ml-0.5 μ g/ml.




Figure 1: Western blot of nonreduced(-) and reduced(+)mouse IgG1 (20 ng/lane), using 0.2 μ g/ml of Clone: RM106. This antibody reacts to nonreduced IgG1 (~150 kDa) stronger than the reduced IgG1 for Clone: RM106 (~50 kDa).



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: ELISA of mouse immunoglobulins shows Clone: RM106 reacts to the Fc region of mouse IgG1; no cross reactivity with IgG2a, IgG3, IgM, IgA, IgE, human IgG, or rat IgG. The plate was coated with 50 ng/well of different immunoglobulins. 200 ng/mL, 50 ng/mL, or 10 ng/mL of Clone: RM106 was used as the primary antibody. An alkaline phosphatase conjugated anti- rabbit IgG as the secondary antibody.




Figure 3: A titer ELISA of mouse IgG1. The plate was coated with different amount of mouse IgG1. A serial dilution of Clone: RM106 was used as the primary antibody. An alkaline phosphatase conjugated anti-rabbit IgG as the secondary antibody.