

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

10-9514-B: Biotinylated Recombinant Goat Monoclonal Antibody to Mouse IgG Fab (Clone: RMG05)(Discontinued)

Clonality: Monoclonal Clone Name: RMG05

Application: IP,ICC,IHC,FACS,ELISA

Reactivity: Mouse
Conjugate: Biotin
Format: Purified
Isotype: Goat IgG
Immunogen Information: Mouse IgG

Product Info

Amount: 50 µg

Purification: Protein G affinity purified from an animal origin–free culture supernatant **Content:** 1 mg/ml in 50% Glycerol/PBS with 1% BSA and 0.09% sodium azide

Storage condition : Store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note

Clone RMG05 antibody reacts to the Fab region of Mouse IgG, including IgG1, IgG2a, IgG2b, and IgG3. No cross reactivity with IgM, IgA, IgE, human IgG, rat IgG, or rabbit IgG. ELISA: 0.05 µg/ml Â- 1 µg/ml; ICC, IHC: 0.5 µg/ml Â- 2 µg/ml.




Figure 1: ELISA of mouse immunoglobulins shows Clone: RMG05 reacts to the Fab region of mouse IgG1, IgG2a, IgG2b, and IgG3; no cross reactivity with IgM, IgA, IgE, human IgG, rat IgG, and rabbit IgG. The plate was coated with 50 ng/well of different immunoglo- bulins. 500 ng/ml, 200 ng/ml, or 50 ng/ml of Clone: RMG05 was used as the primary antibody. An alkaline phosphatase conjugated anti-goat IgG as the secondary antibody.




Figure 2: A titer ELISA of mouse IgG. The plate was coated with different amounts of mouse IgG. A serial dilution of Clone: RMG05 was used as the primary antibody. An alkaline phosphatase conjugated anti-goat IgG as the secondary antibody.