

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

10-9519-B: Biotinylated Recombinant Rabbit Monoclonal Antibody to Human Lambda light chain (Clone: RM127)(Discontinued)

Clonality: Monoclonal Clone Name: RM127

Application: ICC,IHC,FACS,ELISA

Reactivity: Human Conjugate: **Biotin** IGLC1 Gene: Gene ID: 3537 **Uniprot ID:** P0CG04 Format: Purified IGLC1 **Alternative Name:** Isotype: Rabbit IgG Immunogen Information: Human IgG

Product Info

Amount: 50 μg

Purification: Protein A affinity purified from an animal origin-free culture supernatant

Content: 1 mg/ml in 50% Glycerol/PBS with 1% BSA and 0.09% sodium azide

Storage condition : Store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note

Clone RM127 reacts to the lambda light chain of human immunoglobulins. No cross reactivity with kappa light chain, mouse IgG, rat IgG, or goat IgG ELISA: 0.05 $\hat{A}\mu g/ml$ $\hat{A}-$ 0.2 $\hat{A}\mu g/ml$; Immunocytochemistry (ICC): 0.5 $\hat{A}\mu g/ml$ -2 $\hat{A}\mu g/ml$. Immunohistochemistry (IHC): 0.5 $\hat{A}\mu g/ml$ -2 $\hat{A}\mu g/ml$.




Figure 1: Immunohistochemistry of Human Tonsil using Anti- Lambda Light Chain antibody Clone: RM127.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: ELISA of human immunoglobulins shows Clone: RM127 reacts to the Lambda light chain of human immuno-globulins. No cross reactivity with the Kappa light chain, mouse IgG, rat IgG, or goat IgG. The plate was coated with 50 ng/well of different immuno-globulins. 200 ng/mL or 50 ng/mL of Clone: RM127 was used as the primary antibody. An alkaline phos-phatase conjugated anti-rabbit IgG as the secondary antibody.




Figure 3: A titer ELISA using Clone: RM127. The plate was coated with different amounts of human IgG3 Lambda . A serial dilution of Clone: RM127 was used as the primary antibody. An alkaline phosphatase conjugated anti-rabbit IgG as the secondary antibody.