

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

10-9525-B: Biotinylated Recombinant Rabbit Monoclonal Antibody to Human IgD (Clone: RM123)(Discontinued)

Clonality: Monoclonal Clone Name: RM123

Application : ICC,IHC,FACS,ELISA

Reactivity: Human Conjugate: Biotin Gene: **IGHD** Gene ID: 3495 **Uniprot ID:** P01880 Format: Purified **Alternative Name: IGHD** Isotype: Rabbit IgG Immunogen Information: Human IgD

Product Info

Amount: $50 \mu g$

Purification : Protein A affinity purified from an animal origin-free culture supernatant **Content :** 1 mg/ml in 50% Glycerol/PBS with 1% BSA and 0.09% sodium azide

Storage condition : Store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note

Clone RM123 reacts to human IgD. No cross reactivity with human IgG, IgM, IgE, or IgA. ELISA: 25ng/well \tilde{A} [\hat{a} [] 200ng/well (for Capture); 0.01 \tilde{A} [\hat{A} µg/ml \tilde{A} [\hat{a}] 0.1 \tilde{A} [\hat{A} µg/ml (for Detection); Immunocytochemistry (ICC): 0.5 \tilde{A} [\hat{A} µg/ml-2 \tilde{A} [\hat{A} µg/ml] Immunohistochemistry (IHC): 0.5 \tilde{A} [\hat{A} µg/ml-2 \tilde{A} [\hat{A} µg/ml.




Figure 1: Immunohistochemistry of Human Lymphoid Tissue using Anti-Human IgD antibody Clone: RM123.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Sandwich ELISA using Clone: RM123 as the capture antibody (100 ng/well), and Biotinylated anti-human light chains (Kappa+ Lambda) antibody Clone: RM129 as the detection antibody, followed by an alkaline phosphatase conjugated streptavidin.




Figure 3: ELISA of human immunoglobulins shows Clone: RM123 reacts to IgD from human plasma and IgD Kappa from human myeloma. No cross reactivity with human IgG, IgM, IgA, or IgE. The plate was coated with 50 ng/well of different immunoglobulins. 200 ng/mL, 50 ng/mL, or 10 ng/mL of Clone: RM123 was used as the primary antibody. An alkaline phosphatase conjugated anti-rabbit IgG as the secondary antibody.




Figure 4: A titer ELISA using Clone: RM123. The plate was coated with different amounts of human IgD (from plasma). A serial dilution of Clone: RM123 was used as the primary antibody. An alkaline phosphatase conjugated anti-rabbit IgG as the secondary antibody.