

## 10-6601: Mouse Monoclonal Antibody to AIM2 (Clone: 812CT9.2.2.1)(Discontinued)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	812CT9.2.2.1
<b>Application :</b>	WB
<b>Reactivity :</b>	Human
<b>Gene :</b>	AIM2
<b>Gene ID :</b>	9447
<b>Uniprot ID :</b>	O14862
<b>Format :</b>	Purified
<b>Alternative Name :</b>	Interferon-inducible protein AIM2, Absent in melanoma 2, AIM2
<b>Isotype :</b>	Mouse IgM
<b>Immunogen Information :</b>	Synthetic Peptide

### Description

Tumor suppressor which may act by repressing NF-kappa-B transcriptional activity.

### Product Info

<b>Amount :</b>	80 µl / 400 µl
<b>Content :</b>	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Euglobin precipitation followed by dialysis against PBS.
<b>Storage condition :</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term store at -20°C in small aliquots to prevent freeze-thaw cycles.

### Application Note

WB~1:1000




Figure 1: Western blot analysis of AIM2 Antibody (10-6601) with NCI-H292 cell line. AIM2 Antibody was diluted at 1:1000. A goat anti-mouse IgG H&L(HRP) at 1:3000 dilution was used as the secondary antibody. Lysate at 20ug.




Figure 2: Western blot analysis of AIM2 Antibody (10-6601) with NCI-H292 cell line. AIM2 Antibody was diluted at 1:1000. A goat anti-mouse IgG H&L(HRP) at 1:3000 dilution was used as the secondary antibody. Lysate at 20 $\mu$ g.