

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

10-6616: Mouse Monoclonal Antibody to MAFK (Clone: 1328CT786.105.125)(Discontinued)

Clonality: Monoclonal

Clone Name: 1328CT786.105.125

Application: WB,FACS
Reactivity: Human
Gene: MAFK
Gene ID: 7975
Uniprot ID: O60675
Format: Purified

Alternative Name: Transcription factor MafK, Erythroid transcription factor NF-E2 p18 subunit, MAFK

Isotype : Mouse IgG1,Kappa **Immunogen Information :** Recombinant Protein

Description

Since they lack a putative transactivation domain, the small Mafs behave as transcriptional repressors when they dimerize among themselves. However, they seem to serve as transcriptional activators by dimerizing with other (usually larger) basic-zipper proteins and recruiting them to specific DNA-binding sites. Small Maf proteins heterodimerize with Fos and may act as competitive repressors of the NF-E2 transcription factor.

Product Info

Amount : 80 μl / 400 μl

Purification: Protein G Chromatography

Content: Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

Storage condition:

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term store at -20°C in small aliquots to

prevent freeze-thaw cycles.

Application Note

FACS~1:25|| WB~1:1000




Figure 1: Flow cytometric analysis of Hela cells using MAFK Antibody (green) (10-6616) compared to an isotype control of mouse IgG1 (blue). MAFK Antibody was diluted at 1:25 dilution. An Alexa Fluor® 488 goat anti-mouse IgG at 1:400 dilution was used as the secondary antibody.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Western blot analysis of MAFK Antibody (10-6616) lysate from Jurkat cell line . MAFK Antibody was diluted at 1:1000. A goat anti-mouse IgG H&L (HRP) at 1:3000 dilution was used as the secondary antibody. Lysate at $35\mu g$.