

## 10-6616: Mouse Monoclonal Antibody to MAFK (Clone: 1328CT786.105.125)(Discontinued)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	1328CT786.105.125
<b>Application :</b>	WB,FACS
<b>Reactivity :</b>	Human
<b>Gene :</b>	MAFK
<b>Gene ID :</b>	7975
<b>Uniprot ID :</b>	O60675
<b>Format :</b>	Purified
<b>Alternative Name :</b>	Transcription factor MafK, Erythroid transcription factor NF-E2 p18 subunit, MAFK
<b>Isotype :</b>	Mouse IgG1,Kappa
<b>Immunogen Information :</b>	Recombinant Protein

### Description

Since they lack a putative transactivation domain, the small Mafs behave as transcriptional repressors when they dimerize among themselves. However, they seem to serve as transcriptional activators by dimerizing with other (usually larger) basic-zipper proteins and recruiting them to specific DNA-binding sites. Small Maf proteins heterodimerize with Fos and may act as competitive repressors of the NF-E2 transcription factor.

### Product Info

<b>Amount :</b>	80 µl / 400 µl
<b>Purification :</b>	Protein G Chromatography
<b>Content :</b>	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
<b>Storage condition :</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term store at -20°C in small aliquots to prevent freeze-thaw cycles.

### Application Note

FACS~1:25|| WB~1:1000




Figure 1: Flow cytometric analysis of HeLa cells using MAFK Antibody (green) (10-6616) compared to an isotype control of mouse IgG1 (blue). MAFK Antibody was diluted at 1:25 dilution. An Alexa Fluor® 488 goat anti-mouse IgG at 1:400 dilution was used as the secondary antibody.




Figure 2: Western blot analysis of MAFK Antibody (10-6616) lysate from Jurkat cell line . MAFK Antibody was diluted at 1:1000. A goat anti-mouse IgG H&L (HRP) at 1:3000 dilution was used as the secondary antibody. Lysate at 35 $\mu$ g.