

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

10-3010-F: Monoclonal Antibody to TLR4 / CD284 (Clone: ABM19C4) FITC Conjugated

Clonality: Monoclonal ABM19C4 Clone Name: Application: **FACS** Reactivity: Human Conjugate: **FITC** TLR4 Gene: Gene ID: 7099 **Uniprot ID:** O00206 Format: Purified TLR4 **Alternative Name:**

Isotype: Mouse IgG2b Kappa

Immunogen Information: A partial length recombinant TLR4 protein (amino acids 45-320) was used as the immunogen for

this antibody.

Description

TLR4 (Toll-like receptor-4) is a member of the Toll-like receptor (TLR) family which plays a fundamental role in pathogen recognition and activation of innate immunity. It primarily mediates cellular signaling induced by Gram negative bacteria. TLR4 is highly specific for LPS and it is associated with CD14, a co-receptor for LPS. The downstream signaling pathway of TLR4 leads to the activation of NF-kappaB through myeloid differentiation protein (MyD88) and IL-1 receptor-associated kinase in various cell types. Keratinocytes express TLR4 at the mRNA and protein levels whereas TLR4 are also present in the normal human epidermis in vivo and their expression is regulated by microbial components. TLR4 is expressed on CD4+ T cells as well, the functional significance of which is unclear.

Product Info

Amount : $25 \mu g / 100 \mu g$

Purification: Protein G Chromatography

Content: 0.2 mg/ml in Tris buffer and 0.05% sodium azide. Sodium azide is highly toxic.

Storage condition : Store the antibody at 4°C, stable for 6 months.

Application Note

FACS analysis: 2-4 µg/10^6



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig-1 :Flow analysis of human TLR4 antibody (Clone:ABM19C4): Cell surface staining of PBMC, Red represents hTLR4 FITC, Green represents Isotype control mouse IgG2b FITC. $0.5~\mu g$ antibody was used.