

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

10-12502: Mouse Monoclonal Antibody to Bcl2(Clone :BS94)

Clonality: Monoclonal
Clone Name: BS94
Application: IHC
Reactivity: Human
Gene: BCL2
Gene ID: 596
Uniprot ID: P10415

Product Info

Amount: 0.1 ml / 0.5 ml

Content: TRIS with 0.03% sodium azide, pH7.2

Storage condition: Store at 4°C

Application Note

Immunohistochemical Analysis:-1:200



Figure-1: Tonsil (Figure-1) and lymph node sections (Figure-2, figure-3) have been stained using BCL2 antibody (Clone: BS94) with 1:200 dilution. All peripheral lymphocytes should be labelled and the germinal center should be negatively stained from tonsil sections. Only the few scattered T-cells should be stained (Figure-2, figure-3). Mantle cell lymphoma stained strongly with membranous staining pattern.




Figure-2: Tonsil (Figure-1) and lymph node sections (Figure-2, figure-3) have been stained using BCL2 antibody (Clone: BS94) with 1:200 dilution. All peripheral lymphocytes should be labelled and the germinal center should be negatively stained from tonsil sections. Only the few scattered T-cells should be stained (Figure-2, figure-3). Mantle cell lymphoma stained strongly with membranous staining pattern.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com



Figure-3: Tonsil (Figure-1) and lymph node sections (Figure-2, figure-3) have been stained using BCL2 antibody (Clone: BS94) with 1:200 dilution. All peripheral lymphocytes should be labelled and the germinal center should be negatively stained from tonsil sections. Only the few scattered T-cells should be stained (Figure-2, figure-3). Mantle cell lymphoma stained strongly with membranous staining pattern.