

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 39-1015: Anti-CD4 Monoclonal Antibody (Clone: CA-4)

Clonality: Monoclonal

Clone Name: CA-4
Application: IHC-F
Reactivity: Human
Gene: Cd4
Gene ID: 24932
Uniprot ID: P05540

Alternative Name: T-cell surface glycoprotein CD4; T-cell surface antigen T4/Leu-3; W3/25 antigen; CD4; Cd4

**Isotype:** Mouse IgG1

Immunogen Information: CD4-transfected mouse T-cell hybridoma, 3DT, followed by CD4+ human T-cell CEM cells.

## **Description**

CD stands for "cluster of differentiation"; the number that follows is arbitrarily assigned. In the full designation the cell type and nature and molecular weight of the antigen are given in brackets; for CD4, this is as follows: [T,gp55]. CD4 is present on a subset of T lymphocytes("helper/inducer" T cells) and is also expressed at a lower level on monocytes, tissue macrophages and granulocytes. The antigen is involved in binding to MHC class II molecules. The intracellular domain of the antigen is associated with p56lck protein tyrosine kinase.

## **Product Info**

Amount :  $100 \mu g/vial$ Purification : Ascites

Content: Mouse ascites fluid, 1.2% sodium acetate, 2mg BSA, with 0.01mg NaN3 as preservative.

Reconstitute: Add 1ml of PBS buffer will yield a concentration of 100ug/ml.

Storage condition:

At -20?C for one year. After reconstitution, at 4?C for one month. It can also be aliquotted and

stored frozen at -20?C for a longer time. Avoid repeated freezing and thawing.

## **Application Note**

Immunohistochemistry(Frozen Section): 1μg/ml