

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 39-1043: Anti-HSP70 Monoclonal Antibody (Clone: BRM-22)

Clone Name: BRM-22
Application: WB,IHC-P,IHC-F

Reactivity: Human
Gene: Hspa1b
Gene ID: 15511
Uniprot ID: P17879

Alternative Name: Heat shock 70 kDa protein 1B; Heat shock 70 kDa protein 1; HSP70.1; Hspa1b; Hcp70.1,

Hsp70-1, Hsp70a1, Hspa1

**Isotype:** Mouse IgG1

**Immunogen Information:** HSP70 isolated from bovine brain.

## **Description**

Heat-shock proteins, or stress proteins, are expressed in response to heat shock and a variety of other stress stimuli including oxidative free radicals and toxic metal ions. Sargent et al. identified a duplicated HSP70 locus in the class III region of the major histocompatibility complex on 6p21.3. A duplicated locus encoding the major heat shock-induced protein HSP70 is located in the major histocompatibility complex(MHC) class III region 92 kilobases(kb) telomeric to the C2 gene. The 70-kd mammalian heat shock proteins are structurally and functionally related to the uncoating protein that releases clathrin triskelia from coated vesicles.

## **Product Info**

**Amount :** 100 μg/vial **Purification :** Ascites

Content: Mouse ascites fluid, 1.2% sodium acetate, 2mg BSA, with 0.01mg NaN3 as preservative.

Reconstitute: Add 1ml of PBS buffer will yield a concentration of 100ug/ml.

Storage condition:

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and

stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

## **Application Note**

 $We stern \qquad blot \qquad : \qquad 0.5 \tilde{\mathbb{A}} \square \hat{\mathbb{A}} \sqrt[4]{g/ml}; \qquad Immunohistochemistry (Paraffin-embedded \qquad Section) \qquad : \qquad 0.5 - 1 \tilde{\mathbb{A}} \square \hat{\mathbb{A}} \sqrt[4]{g/ml};$ 

Immunohistochemistry(Frozen Section): 0.5-1Ã\(\hat{A}\)\(\hat{4}\)g/ml