w abeomics

39-1046: Anti-MAP Kinase, Activated (Diphosphorylated ERK-1&2) Monoclonal Antibody (Clone: MAPK-YT)

Clonality :	Monoclonal
Clone Name :	MAPK-YT
Application :	ICC,WB
Reactivity :	Human
Gene :	Mapk3
Gene ID :	50689
Uniprot ID :	P21708
Alternative Name :	Mitogen-activated protein kinase 3; MAP kinase 3; MAPK 3; 2.7.11.24; ERT2; Extracellular signal-regulated kinase 1; ERK-1; Insulin-stimulated MAP2 kinase; MAP kinase isoform p44; p44-MAPK; MNK1; Microtubule-associated protein 2 kinase; p44-ERK1; Mapk3; Erk1, Prkm3
Isotype :	Mouse IgG1
Immunogen Information :	A synthetic peptide containing 11 amino acids, HTGFLTpEYpVAT, corresponding to the phosphorylated form of ERK-activation loop conjugated to KLH.

Description

In mammalian cells, a variety of extracellular stimuli generate intracellular signals that converge on a limited number of socalled mitogen-activated protein(MAP) kinase pathways. The central core of each MAP kinase(MAPK) pathway is a conserved cascade of 3 protein kinases: an activated MAPK kinase kinase(MAPKKK) phosphorylates and activates a specific MAPK kinase(MAPKK), which then activates a specific MAPK. Mek1/2 MAPK kinases are essential for mammalian development, homeostasis, and Raf-induced hyperplasia. Germline mutations in genes within the MAPK pathway cause cardio-faciocutaneous syndrome.

Product Info

Amount :	100 µg/vial
Purification :	Ascites
Content :	Mouse ascites fluid, 1.2% sodium acetate, 2mg BSA, with 0.01mg NaN3 as preservative. Reconstitute : Add 1ml of PBS buffer will yield a concentration of 100ug/ml.
Storage condition :	At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Application Note

Western blot : $0.25-0.5\mu$ g/ml; Immunohistochemistry(Paraffin-embedded Section) : $0.4-1\mu$ g/ml; Immunocytochemistry : 1μ g/ml