

### 39-1085: Anti-PCK-26 Monoclonal Antibody (Clone: PCK-26)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	PCK-26
<b>Application :</b>	WB,IHC-P,IHC-F
<b>Reactivity :</b>	Human
<b>Gene :</b>	KRT76
<b>Gene ID :</b>	51350
<b>Uniprot ID :</b>	Q01546
<b>Alternative Name :</b>	Cytokeratin-2P, Keratin-76, Type-II keratin Kb9, KRT2B, KRT2P
<b>Isotype :</b>	Mouse IgG1
<b>Immunogen Information :</b>	Cytokeratin from human epidermis.

#### Description

Monoclonal anti cytokeratins are specific markers of epithelial cell differentiation and have been widely used as tools in tumor identification and classification. Monoclonal Anti Pan Cytokeratin(mixture) is a broadly reactive reagent, which recognizes epitopes present in most human epithelial tissues.

#### Product Info

<b>Amount :</b>	100 µg/vial
<b>Purification :</b>	Ascites
<b>Content :</b>	Mouse ascites fluid, 1.2% sodium acetate, 2mg BSA, with 0.01mg NaN3 as preservative. Reconstitute : Add 1ml of PBS buffer will yield a concentration of 100ug/ml.
<b>Storage condition :</b>	At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

#### Application Note

Western blot : 0.5-1½g/ml; Immunohistochemistry(Paraffin-embedded Section) : 1-2½g/ml; Immunohistochemistry(Frozen Section) : 1-2½g/ml