

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 39-2002: Anti-IL3 Polyclonal Antibody

Clonality: Polyclonal
Application: WB,ELISA
Reactivity: Rat
Gene: II3
Uniprot ID: P04823
Format: Lyophilized

Alternative Name: Interleukin-3; IL-3; Hematopoietic growth factor; Mast cell growth factor; MCGF; Multipotential

colony-stimulating factor; P-cell-stimulating factor; II3; II-3

**Isotype:** Rabbit IgG

Immunogen Information: E. coli-derived rat IL3 recombinant protein (Position: S28-C166). Rat IL3 shares 32% and 55%

amino acid (aa) sequence identity with human and mouse IL3, respectively.

## **Description**

Interleukin 3, also known as IL-3, is a protein that in humans is encoded by the IL3 gene. It is mapped to 5q31.1. IL-3 is an interleukin, a type of biological signal (cytokine) that can improve the body's natural response to disease as part of the immune system. It acts by binding to the interleukin-3 receptor. IL-3 stimulates the differentiation of multipotent hematopoietic stem cells into myeloid progenitor cells or, with the addition of IL-7, into lymphoid progenitor cells. In addition, IL-3 stimulates proliferation of all cells in the myeloid lineage (granulocytes, monocytes, and dendritic cells), in conjunction with other cytokines, e.g., Erythropoietin (EPO), Granulocyte macrophage colony-stimulating factor (GM-CSF), and IL-6. IL-3 is secreted by basophils and activated T cells to support growth and differentiation of T cells from the bone marrow in an immune response.

## **Product Info**

Amount: 100 µg/vial

**Purification:** Immunogen affinity purified.

Content: Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3. Reconstitute: Add

0.2ml of distilled water will yield a concentration of 500ug/ml.

Storage condition:

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and

stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

## **Application Note**

Western blot :  $0.1-0.5\tilde{A} \square \hat{A}^{1/4}q/ml$ ; ELISA :  $0.1-0.5\tilde{A} \square \hat{A}^{1/4}q/ml$ 




Figure 1: Anti-IL-3 antibody(39-2002) . Western blotting : Lanes: Anti IL-3(39-2002) at 0.5 $\mu$ ml. WB: Recombinant Rat IL-3 Protein 0.5 $\mu$ ml. Predicted band size: 15 kDa.Observed band size: 15 kDa.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Anti- IL-3 antibody(39-2002). Western blotting : - Lanes: Anti IL-3(39-2002) at 0.5ug/ml, Lane 1: PC-12 Whole Cell Lysate at 40ug, Lane 2: NRK Whole Cell Lysate at 40ug, Lane 3: RH35 Whole Cell Lysate at 40ug; Predicted band size: 17 kDa: Observed band size: 37 kDa.