

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

39-2014: Anti-Alkaline Phosphatase Polyclonal Antibody

Clonality: Polyclonal

Application:WBReactivity:RatGene:AlplGene ID:25586Uniprot ID:P08289Format:Lyophilized

Alkaline phosphatase, tissue-nonspecific isozyme; AP-TNAP; TNSALP; 3.1.3.1; Alkaline

phosphatase liver/bone/kidney isozyme; Alpl

Isotype: Rabbit IgG

A synthetic peptide corresponding to a sequence at the N-terminus of rat Alkaline

Immunogen Information: Phosphatase(18-32aa, FVPEKEKDPSYWRQQ), different from the related mouse sequence by

one amino acid.

Description

Alkaline phosphatase, tissue-nonspecific isozyme is an enzyme that in humans is encoded by the ALPL gene. There are at least four distinct but related alkaline phosphatases: intestinal, placental, placental-like, and liver/bone/kidney(tissue non-specific). The first three are located together on chromosome 2 while the tissue non-specific form is located on chromosome 1. Liver/bone/kidney alkaline phosphatase, also known as tissue-nonspecific alkaline phosphatase, acts physiologically as a lipid-anchored phosphoethanolamine(PEA) and pyridoxal-5-prime-phosphate(PLP) ectophosphatase Mice that lack a functional form of this enzyme show normal skeletal development. This enzyme has been linked directly to a disorder known as hypophosphatasia, a disorder that is characterized by hypercalcemia and includes skeletal defects.

Product Info

Amount: 100 μg/vial

Purification: Immunogen affinity purified.

Content: Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.

Reconstitute: Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Storage condition:

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and

stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Application Note

Western blot : 0.1-0.5̸¼g/ml




Figure 1: All Lanes: Anti ALPL(39-2014) at 0.5ug/ml. Lane 1: Rat Cardiac Muscle Tissue Lysate at 50ug, Lane 2: Rat Brain Tissue Lysate at 50ug. Predicted band size: 57 kDa. Observed band size: 57 kDa.