

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 39-2016: Anti-Angiopoietin 2 Polyclonal Antibody

**Clonality:** Polyclonal

Application: WB
Reactivity: Human
Gene: ANGPT2
Gene ID: 285
Uniprot ID: O15123
Format: Lyophilized

**Alternative Name :** Angiopoietin-2; ANG-2; ANGPT2

**Isotype:** Rabbit IgG

Immunogen Information: A synthetic peptide corresponding to a sequence at the C-terminus of human Angiopoietin

2(454-468aa NLNGMYYPQRQNTNK).

## **Description**

Angiopoietin 1 and Angiopoietin 2 are important for development of the endothelium, by regulating tyrosine phosphorylation of the membrane receptor Tie 2. Angiopoietin 2 is only 60% homologous with Angiopoietin 1. Angiopoietin-2 is a naturally occurring antagonist of angiopoietin-1 that competes for binding to the TIE2 receptor and blocks ANGPT1-induced TIE2 autophosphorylation. Angiopoietin 1 binding to Tie 2 causes phosphorylation of the receptor. Angiopoietin 2 competes for this binding, and thus blocks receptor phosphorylation. Angiopoietin 2 expression occurs at sites of vascular remodelling: dorsal aorta and major aortic branches, ovary, placenta and uterus.

## **Product Info**

Amount: 100 μg/vial

**Purification:** Immunogen affinity purified.

Content: Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.

Reconstitute: Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Storage condition:

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and

stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

## **Application Note**

Western blot :  $0.1-0.5\tilde{A} \square \hat{A}^{1/4}g/ml$ ;




Figure 1: Anti-Angiopoietin 2 antibody(39-2016). Western blotting: Lane 1: Recombinant Human ANG2 Protein 10ng, Lane 2: Recombinant Human ANG2 Protein 5ng. Lane 3: Recombinant Human ANG2 Protein 2.5ng.