

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 39-2023: Polyclonal Antibody to Anti-ATF6 Antibody(Discontinued)

Clonality: Polyclonal
Reactivity: Human
Gene: ATF6
Gene ID: 22926
Uniprot ID: P18850

Alternative Name:

Cyclic AMP-dependent transcription factor ATF-6 alpha; cAMP-dependent transcription factor

ATF-6 alpha; Activating transcription factor 6 alpha; ATF6-alpha

**Isotype:** Rabbit IgG

A synthetic peptide corresponding to a sequence in the middle region of human

Immunogen Information: ATF6(390-405aa LNYGPMSMLEQDSRRM), different from the related mouse and rat sequences

by two amino acids.

## **Description**

ATF6, a member of the leucine zipper protein family, is an endoplasmic reticulum(ER) stress-regulated transmembrane transcription factor that activates the transcription of ER molecules. ATF6 gene is mapped to chromosome 1q23.3. ATF6 can constitutively induce the promoter of glucose-regulated protein(grp) genes through activation of the endoplasmic reticulum(ER) stress element(ERSE).

## **Product Info**

Amount: 100 μg/vial

**Purification:** Immunogen affinity purified.

Content: Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.

Reconstitute: Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Storage condition:

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and

stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

## **Application Note**

Western blot :  $0.1\text{-}0.5\mu\text{g/ml}$ 

130KD — 100KD — — 70KD — 55KD —

25KD -

Figure 1: Anti-ATF6 antibody(39-2023). Western blotting: Lanes: Anti ATF6 39-2023 at 0.5ug/ml. WB: A549 Whole Cell Lysate at 40ug. Predicted band size: 75 kDa. Observed band size: 90 kDa.