

39-2054: Anti-NMDAR2A Polyclonal Antibody

| | |
|---------------------------|--|
| Clonality : | Polyclonal |
| Application : | WB |
| Reactivity : | Human |
| Gene : | GRIN2A |
| Gene ID : | 2903 |
| Uniprot ID : | Q12879 |
| Alternative Name : | Glutamate receptor ionotropic, NMDA 2A; GluN2A; Glutamate [NMDA] receptor subunit epsilon-1; N-methyl D-aspartate receptor subtype 2A; NMDAR2A; NR2A; hNR2A; GRIN2A; NMDAR2A |
| Isotype : | Rabbit IgG |

Description

GRIN2A is also known as N-methyl-D-aspartate receptor channel, subunit epsilon-1(NMDAR2A). This gene encodes a member of the glutamate-gated ion channel protein family. The encoded protein is an N-methyl-D-aspartate (NMDA) receptor subunit. NMDA receptors are both ligand-gated and voltage-dependent, and are involved in long-term potentiation, an activity-dependent increase in the efficiency of synaptic transmission thought to underlie certain kinds of memory and learning. These receptors are permeable to calcium ions, and activation results in a calcium influx into post-synaptic cells, which results in the activation of several signaling cascades. Disruption of this gene is associated with focal epilepsy and speech disorder with or without mental retardation. Alternative splicing results in multiple transcript variants.

Product Info

| | |
|----------------------------|---|
| Amount : | 100 µg/vial |
| Purification : | Immunogen affinity purified. |
| Content : | Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg Na ₃ . Reconstitute : Add 0.2ml of distilled water will yield a concentration of 500ug/ml. |
| Storage condition : | At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing. |

Application Note

Western blot : 0.1-0.5µg/ml; Immunohistochemistry(Paraffin-embedded Section) : 0.5-1µg/ml




Figure 1: Anti-NMDAR2A antibody(39-2054). Western blotting: Lanes: Anti NMDAR2A(39-2054) at 0.5ug/ml, Lane 1: Rat Brain Tissue Lysate at 50ug, Lane 2: Mouse Brain Tissue Lysate at 50ug. Predicted band size: 165 kDa. Observed band size: 165 kDa.




Figure 2: Anti-NMDAR2A antibody(39-2054). IHC(P): Rat Brain Tissue.