

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

39-2155: Anti-TNF alpha Polyclonal Antibody

Clonality: Polyclonal
Application: WB
Reactivity: Rat
Gene: Tnf

 Gene ID :
 103694380

 Uniprot ID :
 P16599

 Format :
 Lyophilized

Tumor necrosis factor; Cachectin; TNF-alpha; Tumor necrosis factor ligand superfamily member 2;

Alternative Name: TNF-a; Tumor necrosis factor, membrane form; N-terminal fragment; NTF; Intracellular domain 1;

ICD1

Isotype: Rabbit IgG

Immunogen Information: E.coli-derived rat TNF alpha recombinant protein (Position: D89-L235). Rat TNF alpha shares 95%

amino acid (aa) sequence identity with mouse TNF alpha.

Description

TNF alpha(Tumor Necrosis Factor alpha) gene encodes a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor (TNF) superfamily. This cytokine is mainly secreted by macrophages. It can bind to, and thus functions through its receptors TNFRSF1A/TNFR1 and TNFRSF1B/TNFBR. This cytokine is involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. This cytokine has been implicated in a variety of diseases, including autoimmune diseases, insulin resistance, and cancer. Knockout studies in mice also suggested the neuroprotective function of this cytokine.

Product Info

Amount: $100 \mu g/vial$

Purification: Immunogen affinity purified.

Content: Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3. Reconstitute : Add

0.2ml of distilled water will yield a concentration of 500ug/ml.

Storage condition:

At -20?C for one year. After reconstitution, at 4?C for one month. It can also be aliquotted and

stored frozen at -20?C for a longer time. Avoid repeated freezing and thawing.

Application Note

Western blot: 0.1-0.5Î1/4g/ml




Figure 1: Anti-TNF alpha antibody(39-2155). Lanes: Anti-TNF alpha(39-2155) at 0.5ug/ml. Lane 1: PC-12 Whole Cell Lysate at 40ug, Lane 2: Rat Spleen Tissue Lysate at 40ug, Lane 3: Rat Brain Tissue Lysate at 40ug, Lane 4: Rat Kidney Tissue Lysate at 40ug, Lane 5: Rat Liver Tissue Lysate at 40ug. Predicted band size: 26 kDa. Observed band size: 26 kDa.