

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

39-2158: Anti-IL2 Polyclonal Antibody

Clonality: Polyclonal

 Application :
 WB

 Reactivity :
 Rat

 Gene :
 II2

 Gene ID :
 116562

 Uniprot ID :
 P17108

 Format :
 Lyophilized

Alternative Name: Interleukin-2; IL-2; T-cell growth factor; TCGF; II2; II-2

Isotype: Rabbit IgG

Immunogen Information: E.coli-derived rat IL2 recombinant protein (Position: A21-Q155). Rat IL2 shares 66% and 80%

amino acid (aa) sequence identity with human and mouse IL2, respectively.

Description

Interleukin-2 (IL2), formerly referred to as T-cell growth factor, is a powerfully immunoregulatory lymphokine that is produced by lectin- or antigen-activated T cells. It is produced not only by mature T lymphocytes on stimulation but also constitutively by certain T-cell lymphoma cell lines. The lymphokine interleukin-2 (IL-2) is responsible for autocrine cell cycle progression and regulation of immune responses. IL-2 expression in mature thymocytes and T cells has been found to be tightly controlled by monoallelic expression. IL-2 can act as a growth hormone for both B and T lymphocytes. This gene for interleukin 2 (IL2) is assigned to chromosome 4. IL-2 is a 15.5KDa glycoprotein, consisting of 153 amino acids in precursor form and 133 amino acids in mature form.

Product Info

Amount: 100 μg/vial

Purification: Immunogen affinity purified.

Content: Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3. Reconstitute : Add

0.2ml of distilled water will yield a concentration of 500ug/ml.

Storage condition:

At -20?C for one year. After reconstitution, at 4?C for one month. It can also be aliquotted and

stored frozen at -20?C for a longer time. Avoid repeated freezing and thawing.

Application Note

Western blot: 0.1-0.5μg/ml




Figure 1: Anti-IL-2 antibody(39-2158). Lanes: Anti-IL-2(39-2158) at 0.5ug/ml. Lane 1: Rat Thymus Tissue Lysate at 40ug, Lane 2: Rat Liver Tissue Lysate at 40ug, Lane 3: NRK Whole Cell Lysate at 40ug, Lane 4: PC12 Whole Cell Lysate at 40ug, Lane 5: RH35 Whole Cell Lysate at 40ug. Predicted band size: 18 kDa. Observed band size: 18 kDa.