

## 10-4137-F: Monoclonal Antibody to CD28 (Clone: CB28) FITC Conjugated

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	CB28
<b>Application :</b>	FACS
<b>Reactivity :</b>	Human
<b>Conjugate :</b>	FITC
<b>Gene :</b>	CD28
<b>Gene ID :</b>	940
<b>Uniprot ID :</b>	P10747
<b>Format :</b>	Purified
<b>Alternative Name :</b>	T-cell-specific surface glycoprotein CD28, TP44, CD28
<b>Isotype :</b>	Mouse IgG1, Kappa
<b>Immunogen Information :</b>	A partial length recombinant protein from CD28 was used as the immunogen for this antibody.

### Description

CD28 is a type I transmembrane protein that binds through its extracellular region to B7 proteins (CD80 and CD86), which are transmembrane proteins expressed on the surface of APCs (Antigen-Presenting Cells) and are up-regulated by inflammatory signals. CD28 mediates signals that promote T lymphocyte differentiation and proliferation, and enhance antibody production of B lymphocytes. Deficiencies in CD28 pathways result in complete T lymphocyte tolerance in vitro and in vivo. Both CD4+ and CD8+ memory T cells need CD28 costimulation to achieve maximal expansion and pathogen clearance. The blockade of the CD28-B7 interaction has been used to down-regulate the activation of the immune system in autoimmune diseases. Based on the expression of the costimulatory molecule CD28 on the surface of CD8+ T cells, two different lymphocyte subgroups have been designated: antigen-primed cytotoxic T cells (CD8+CD28+ T cells) and suppressor T cells (CD8+CD28- T cells). The frequency of CD28+CD8+ T cells and especially the balance between CD8+CD28+ and CD8+CD28- T cells are important in many diseases, including CHB (Chronic Hepatitis B).

### Product Info

<b>Amount :</b>	100 µg
<b>Purification :</b>	Protein G Chromatography
<b>Content :</b>	25 µg in 125 µl/100 µg in 500 µl Tris and 0.05% sodium azide. Sodium azide is highly toxic.
<b>Storage condition :</b>	Store the antibody at 4°C, stable for 6 months.

### Application Note

FACS: 0.2-0.5 µg/10<sup>6</sup> cells




Fig-1: Cell surface Flow analysis of CD28 in PBMC (lymphocytes gated) using 0.2 $\mu$ g/10<sup>6</sup> cells. Green represents FITC conjugated isotype control (ABEOMICS); red represents FITC conjugated anti-hCD28 antibody (10-4137-F).