

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

10-7613: Monoclonal antibody to MSi-2 (Clone: ABM5A48)

Clonality: Monoclonal **Clone Name:** ABM5A48 IHC.FACS.WB Application: Reactivity: Human Gene: MSI2 Gene ID: 124540 **Uniprot ID:** Q96DH6 Format: Purified

Alternative Name: RNA-binding protein Musashi homolog 2, Musashi-2

Isotype: Mouse IgG2b, Kappa

Immunogen Information: A partial length recombinant protein of Msi-2 (amino acid 18-232) was used as the

immunogen for this antibody.

Product Info

Amount : $25 \mu g / 100 \mu g$

Purification: Protein G Chromatography

Content: 25 μg in 50 μl/100 μg in 200 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium

azide is highly toxic.

Storage condition : Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid

repeated freeze and thaw cycles

Application Note

Facs analysis: 2-4 μg/10⁶ Cells, Western blot analysis : 2-4 μg/ml, Immunohistochemical

analysis: 5-10 μg/ml




Figure-1: Intra cellular flow analysis of Msi-2 on HeLa cells using 2 µg/10⁶ cells of Msi-2 antibody (Clone: ABM5A48). Green represents isotype control; red represents anti-Msi-2 antibody. Goat anti-mouse PE conjugate was used as secondary antibody.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure-2: Intra cellular flow analysis of Msi-2 on A431 cells using $2 \mu g/10^6$ cells of Msi-2 antibody (Clone: ABM5A48). Green represents isotype control; red represents anti-Msi-2 antibody. Goat anti-mouse PE conjugate was used as secondary antibody




Figure-3: Western blot analysis of MSI-2. Anti Msi-2 (Clone: ABM5A48) was used at 0.5 μ g/ml in Recombinant and 2 μ g/ml in SKBR3, MCF7 Lysates.




Figure-4:Immunohistochemical analysis of MSI-2 in Adenocarcinoma of Lungs tissue using Anti-Msi-2 (Clone: ABM5A48) at 5 µg/ml.