

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 10-3532: Monoclonal Antibody to mouse M-CSF(Discontinued)

Clonality: Monoclonal
Clone Name: ER-MP58
Application: FACS
Reactivity: Mouse
Gene: Csf1
Gene ID: 12977
Uniprot ID: P07141

Alternative Name: Csfm, Macrophage colony-stimulating factor 1

**Isotype:** Rat IgG2a

## **Description**

The monoclonal antibody ER-MP58 recognizes an antigen present on all mouse macrophage colonystimulating factor (M-CSF) responsive cells in the bone marrow, including the earliest colony-forming myeloid progenitors, as well as by the majority of other myeloid precursors. The antigen is detected on a broad development range of macrophage precursor cells to the monocytic level, but also on granulocytes. Expression is rapidly lost upon maturation beyond the monocytic stage. The antigen disappears in the course of macrophage differentiation. Furthermore the antigen is clearly different from commonly used myeloid markers as Mac-1, F4/80, and Gr-1. The monoclonal antibody ER-MP58 is very useful for the identification of mouse myeloid hematopoietic islands in various organs, and for embryonic studies. Cells committed to the myeloid lineage can be separated from progenitor cells with other differentiation capacities by means of multiparameter cell sorting using monoclonal antibody ER-MP58 in combination with monoclonal antibody ER-MP12 and ER-MP-20.

## **Product Info**

**Amount :** CSF(Discontinued) / 500 μg

Content: 0.5 mg, 0.2 µm filtered antibody solution in PBS, containing 0.1% bovine serum albumin and

0.02% sodium azide.

**Storage condition :** Product should be stored at 4 °C. Under recommended storage conditions, product is stable for

one year.

## **Application Note**

For immunohistology and flow cytometry, dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50.