

32-7026: Recombinant Human Insulin-Like Growth Factor I/IGF-I/IGF1

Gene : IGF1
Gene ID : 3479
Uniprot ID : P05019

Description

Source: E.coli.
MW :7.6kD.

Recombinant Human Insulin-like Growth Factor I is produced by our E.coli expression system and the target gene encoding Gly49-Ala118 is expressed. Insulin-like growth factor I (IGF1) belongs to the family of insulin-like growth factors that are structurally homologous to proinsulin. Mature IGFs are generated by proteolytic processing of inactive precursor proteins, which contains the N- and C-terminal propeptide regions. Mature human IGF-I consisting of 70 amino acids has 94% identity with mouse IGF-I and exhibits cross-species activity. IGF-1 binds IGF-IR, IGF-IIR, and the insulin receptor and plays a key role in cell cycle progression, cell proliferation and tumor progression. IGF-1 expression is regulated by growth hormone. R3 IGF-1 is an 83 amino acid analog of IGF-1 comprising the complete human IGF-1 sequence with the substitution of an Arg (R) for the Glu(E) at position three, hence R3, and a 13 amino acid extension peptide at the N terminus. R3 IGF-1 has been produced with the purpose of increasing biological activity. R3 IGF-1 is significantly more potent than human IGF-I in vitro.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 300mM NaAc, pH 6.5.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : GPETLCGAELVDALQFVCGDRGFYFNKPTGYGSSSRAPQTGIVDECCFRSCDLRRLEMYCAPLK PAKSA

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 Åµg/ml. Dissolve the lyophilized protein in 500mM Acetic Acid. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.05 ng/Åµg (0.5 IEU/Åµg) as determined by LAL test.

Biological Activity : ED50 is greater than 200 ng/ml.