

32-7033: Recombinant Human GM-CSF/CSF2 (P. pastoris)

Gene : CSF2
Gene ID : 1437
Uniprot ID : P04141

Description

Source: P.Pichia.

MW :14.4kD.

Recombinant Human Granulocyte-Macrophage Colony-Stimulating Factor is produced by our Yeast expression system and the target gene encoding Ala18-Glu144 is expressed. GM-CSF was initially characterized as a growth factor that can support the in vitro colony formation of granulocyte macrophage progenitors. It is produced by a number of different cell types (including activated T cells, B cells, macrophages, mast cells, endothelial cells and fibroblasts) in response to cytokine of immune and inflammatory stimuli. Besides granulocyte-macrophage progenitors, GM-CSF is also a growth factor for erythroid, megakaryocyte and eosinophil progenitors. On mature hematopoietic, monocytes/macrophages, and eosinophils, GM-CSF has also been reported to have a functional role on non-hematopoietic cells. It can induce human endothelial cells to migrate and proliferate. Additionally, GM-CSF can also stimulate the proliferation of a number of tumor cell lines, including osteogenic sarcoma, carcinoma and adenocarcinoma cell lines.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 10mM TrisHCl, 4% Mannitol, 1% Sucrose, pH 8.5.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : APARSPSPSTQPWEHVNAIQEARRLLNLSRDTAAEMNETVEVISEMFDLQEPTCLQTRLELYKQGLRGS�TKLK GPLTMMASHYKQHCPTPETSCATQIITFESFKENLKDFLLVIPFDCWEPVQE

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 Åµg/ml. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/Åµg (1 IEU/Åµg) as determined by LAL test.

Biological Activity : ED50 is less than 0.2 ng/ml. Specific Activity of 5.0 x 10⁶ IU/ mg.