

## 32-7037: Recombinant Mouse Fibroblast Growth Factor 2/FGF-2/FGFb (Met1-Ser154)

**Gene :** Fgf2  
**Gene ID :** 14173  
**Uniprot ID :** P15655

### Description

Source: E.coli.  
MW :17.15kD.

Recombinant Mouse Fibroblast growth factor 2 is produced by our E.coli expression system and the target gene encoding Met1-Ser154 is expressed. FGF basic is one of 22 mitogenic proteins of the FGF family, which show 35-60% amino acid conservation. Unlike other FGFs, FGF acidic and basic lack signal peptides and are secreted by an alternate pathway. The 17 kDa mouse sequence has 98% aa identity with rat, and 95% identity with human, bovine, and sheep FGF basic. Binding of FGF to heparin or cell surface HSPG is necessary for binding, dimerization and activation of tyrosine kinase FGF receptors. FGF basic binds other proteins, polysaccharides and lipids with lower affinity. Expression of FGF basic is nearly ubiquitous but disruption of the mouse FGF basic gene gives a relatively mild phenotype, suggesting compensation by other FGF family members. FGF basic modulates such normal processes as angiogenesis, wound healing and tissue repair, embryonic development and differentiation, neuronal function and neural degeneration. Transgenic overexpression of FGF basic results in excessive proliferation and angiogenesis is reminiscent of a variety of pathological conditions.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM PB, 400mM NaCl, pH 7.0.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** MAASGITSLPALPEDGGAFFPPGHFKDPKRLYCKNGGFFLRHPDGRVDGVREKSDPHVKLQLQA  
EERGVSISIKGVCANRYLAMKEDGRLLASKCVTEECFFFERLESNNYNTYRSRKYSWYVALKRTG  
QYKLGSKTGPGQKAILFLPMSAKS

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 Åµg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/Åµg (1 IEU/Åµg) as determined by LAL test.