

32-7043: Recombinant Human Interleukin-4/IL-4 (E. coli)

Gene : IL4
Gene ID : 3565
Uniprot ID : P05112

Description

Source: E.coli.
MW :15.1kD.

Recombinant Human Interleukin-4 is produced by our E.coli expression system and the target gene encoding His25-Ser153 is expressed. Interleukin-4 (IL-4) is a pleiotropic cytokine that regulates diverse T and B cell responses including cell proliferation, survival and gene expression. IL-4 is produced by mast cells, T cells, and bone marrow stromal cells. IL-4 regulates the differentiation of naive CD4+ T cells into helper Th2 cells, characterized by their cytokine-secretion profile that includes secretion of IL-4, IL-5, IL-6, IL-10, and IL-13, which favor a humoral immune response. Another dominant function of IL-4 is the regulation of immunoglobulin class switching to the IgG1 and IgE isotypes. Excessive IL-4 production by Th2 cells has been associated with elevated IgE production and allergic response.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : MHKCDITLQEIIKTLNSLTEQKTLCTELTVTDIFAASKNTEKETFCRAATVLRQFYSHHEKDTRCLG
ATAQQFHRHKQLIRFLKRLDRNLWGLAGLNSCPVKEANQSTLENFLERLKTIMREKYSKCSS

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.

Biological Activity : ED50 is less than 2 ng/ml. Specific Activity of 5.0 x 10⁶ IU/mg.