

## 32-7072: Recombinant Human Interleukin-7/IL-7

**Gene :** IL7  
**Gene ID :** 3574  
**Uniprot ID :** P13232

### Description

Source: E.coli.  
MW :17.5kD.

Recombinant Human Interleukin-7 is produced by our E.coli expression system and the target gene encoding Asp26-His177 is expressed. Human Interleukin 7 (IL-7) is a potent lymphoid cell growth factor stimulating the proliferation of lymphoid progenitors. IL7 can associate with the hepatocyte growth factor (HGF) to form a hybrid cytokine that functions as a pre-pro-B cell growth-stimulating factor. Human IL7 cDNA encodes a 177 amino acid precursor protein containing a 25 amino acid signal peptide and a 152 amino acid mature protein. Human and mouse IL7 share 65% sequence identity in the mature region and both exhibit cross-species activity. IL-7 signals via IL-7 receptor (IL7R) activating multiple pathways including JaK/STAT and PI3K/AKT, which regulate lymphocyte survival, glucose uptake, proliferation, and differentiation. IL-7 is also associated with cytoplasmic IL2-R gamma for signal transduction.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM PB, 250mM NaCl, pH 7.2.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** MDCDIEGKDGGKQYESVLMVSIQQLDSMKEIGSNCLNNEFNFFKRHICDANKEGMFLFRAARKLRQFLKMNSTGDFDLHLLKVSEGTILLNCTGQVKGRKPAALGEAQPTKSLEENKSLKEQKKLNDLCFLKRLQEIKTWNKILMGTKEH

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.