

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-7083: Recombinant Human Ciliary Neurotrophic Factor/CNTF

Gene ID: CNTF
Gene ID: 1270
Uniprot ID: P26441

## **Description**

Source: E.coli. MW:22.93kD.

Recombinant Human CNTF is produced by our E.coli expression system and the target gene encoding Ala2-Met200 is expressed. Ciliary Neurotrophic Factor (CNTF) is a potent survival factor for neurons and oligodendrocytes. CNTF has also been shown to prevent the degeneration of motor axons after axotomy. CNTF is highly conserved across species and exhibits cross-species activities. Human and rat CNTF share approximately 83% homology in their protein sequence. CNTF is structurally related to IL6, IL11, LIF and OSM. All of these four helix bundle cytokines share gp130 as a signal transducing subunit in their receptor complexes. CNTF, like FGF acidic, FGF basic, and PD-ECGF (platelet-derived endothelial cell growth factor), does not possess a signal sequence that would allow secretion of the factor by classical secretion pathways. The mechanism underlying the release of CNTF is unknown.

## **Product Info**

**Amount:**  $10 \mu g / 50 \mu g$ 

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, 1mM Cysteine, pH 7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

**Storage condition :** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: AFTEHSPLTPHRRDLCSRSIWLARKIRSDLTALTESYVKHQGLNKNINLDSADGMPVASTDQWSELTEAERLQE

NLQAYRTFHVLLARLLEDQQVHFTPTEGDFHQAIHTLLLQVAAFAYQIEELMILLEYKIPRNEADGMPINVGDGG

LFEKKLWGLKVLQELSQWTVRSIHDLRFISSHQTGIPARGSHYIANNKKM

## **Application Note**

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\tilde{A} \square \hat{A} \mu g/ml$ . Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin**: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$  (1 IEU/ $\tilde{A} \square \hat{A} \mu g$ ) as determined by LAL test.