

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-7102: Recombinant Human Cornulin (N-6His)

Gene ID: CRNN
Gene ID: 49860
Uniprot ID: Q9UBG3

## **Description**

Source: E.coli. MW:17.45kD.

Recombinant Human Cornulin is produced by our E.coli expression system and the target gene encoding Met1-Ser140 is expressed with a 6His tag at the N-terminus. Cornulin is a member of the fused gene family of molecular chaperones. Human Cornulin contains N-terminus EF-hand domains and Ca2+ binding domains, and two glutamine- and threonine-rich 60 amino acid repeats in its C-terminus. Cornulin involves in the mucosal/epithelial immune response and epidermal differentiation. Cornulin is a survival factor that participates in the clonogenicity of squamous esophageal epithelium cell lines, attenuates deoxycholic acid (DCA)-induced apoptotic cell death and release of calcium. When Cornulin is overexpressed in oral squamous carcinoma cell lines, it regulates negatively cell proliferation by the induction of G1 arrest.

## **Product Info**

**Amount :**  $10 \mu g / 50 \mu g$ 

**Content:** Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

**Storage condition:** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: MGSSHHHHHHSSGLVPRGSHMPQLLQNINGIIEAFRRYARTEGNCTALTRGELKRLLEQEFADVIVKPHDPATV

DEVLRLLDEDHTGTVEFKEFLVLVFKVAQACFKTLSESAEGACGSQESGSLHSGASQELGEGQRSGTEVGRAG

**KGQHYEGSSHRQS** 

## **Application Note**

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\tilde{A} \square \hat{A} \mu g/ml$ . Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$  (1 IEU/ $\tilde{A} \square \hat{A} \mu g$ ) as determined by LAL test.