

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-7120: Recombinant Human Fatty Acid-Binding Protein 7/FABP7/B-FABP (N-6His)

**Gene ID**: PABP7 **Gene ID**: 2173 **Uniprot ID**: O15540

## **Description**

Source: E.coli. MW :17.05kD.

Recombinant Human FABP7 is produced by our E.coli expression system and the target gene encoding Val2-Ala132 is expressed with a 6His tag at the N-terminus. Fatty Acid-Binding Protein 7 (FABP7) is a cytoplasm protein that belongs to the Fatty-acid Binding Protein (FABP) family of calycin superfamily. Fatty acid binding proteins are a family of small, highly conserved, cytoplasmic proteins that bind long-chain fatty acids. FABP7 is predominately expressed in brain and neural tissues. FABP7 is involved in fatty acid uptake and intracellular transport and is important in brain development. FABP7 plays a critical role in the transport of a so far unknown hydrophobic ligand with potential morphogenic activity during CNS development. FABP7 is required for the establishment of the radial glial fiber system in developing brain, a system that is necessary for the migration of immature neurons to establish cortical layers.

## **Product Info**

**Amount :**  $10 \mu g / 50 \mu g$ 

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

**Storage condition :** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: MGSSHHHHHHSSGLVPRGSHMVEAFCATWKLTNSQNFDEYMKALGVGFATRQVGNVTKPTVIIS

QEGDKVVIRTLSTFKNTEISFQLGEEFDETTADDRNCKSVVSLDGDKLVHIQKWDGKETNFVREIKD

GKMVMTLTFGDVVAVRHYEKA

## **Application Note**

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\hat{A}\mu g/ml$ . Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.