

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7123: Recombinant E. coli Formamidopyrimidine-DNA Glycosylase/Fpg (C-6His)(Discontinued)

Gene: mutM **Uniprot ID**: A7ZTI6

Description

Source: E.coli. MW :31.4kD.

Recombinant E.coli Fpg is produced by our E.coli expression system and the target gene encoding Met1-Lys269 is expressed with a 6His tag at the C-terminus. Fpg is a DNA glycosylase that releases damaged bases preferentially from duplex DNA. It has an associated class I AP (apurinic/apyrimidinic) lyase activity. Fpg Cleaves the DNA backbone to generate a single-strand break at the site of the removed base, The C-O-P bond 3' to the apurinic or apyrimidinic site in DNA is broken by a beta-elimination reaction, leaving 3' and 5' phosphoryl groups.

Product Info

Amount: $10 \mu g / 50 \mu g$

Content: Supplied as a 0.2 µm filtered solution of 20mM TrisHCl, 100mM NaCl, 1mM EDTA, 1mM DTT,

50% Glycerol, pH 7.8.

Storage condition : Store at -20°C, stable for 6 months after receipt. Please minimize freeze-thaw cycles.

Amino Acid: MPELPEVETSRRGIEPHLVGATILHAVVRNGRLRWPVSEEIYRLSDQPVLSVQRRAKYLLLELPEG

WIIIHLGMSGSLRILPEELPPEKHDHVDLVMSNGKVLRYTDPRRFGAWLWTKELEGHNVLTHLGPE PLSDDFNGEYLHQKCAKKKTAIKPWLMDNKLVVGVGNIYASESLFAAGIHPDRLASSLSLAECELLA RVIKAVLLRSIEQGGTTLKDFLQSDGKPGYFAQELQVYGRKGEPCRVCGTPIVATKHAQRATFYCR

QCQKLEHHHHHH

Application Note

Endotoxin: Less than 0.1 ng/Âμg (1 IEU/Âμg) as determined by LAL test.