

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7151: Recombinant Human Stratifin/SFN(Discontinued)

Gene ID: 2810 Uniprot ID: P31947

Description

Source: E.coli. MW:27.77kD.

Recombinant Human Stratifin is produced by our E.coli expression system and the target gene encoding Met1-Ser248 is expressed. Stratifin (SFN) belongs to the 14-3-3 family of proteins that act as important regulators of intracelluar signal transduction through their ability to bind specific motifs phosphorylated on serine or threonine. There are at least seven isoforms that have been identified in mammals (beta, gamma, epsilon, sigma, zeta, tau and eta). SFN can detected in many tissues, highly expressed in stratified squamous keratinizing epithelium. SFN is indicated as an epithelial cell marker and serves as a tumor suppressor whose expression can be down regulated by methylation. In addition, SFN plays a key role in maintaining the G2 checkpoint in cells and preventing mitotic death.

Product Info

Amount : $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 μm filtered solution of 20mM PB, 250mM NaCl, 1mM EDTA, 1mM DTT, pH

7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: MERASLIQKAKLAEQAERYEDMAAFMKGAVEKGEELSCEERNLLSVAYKNVVGGQRAAWRVLSSIEQKSNEEG

SEEKGPEVREYREKVETELQGVCDTVLGLLDSHLIKEAGDAESRVFYLKMKGDYYRYLAEVATGDDKKRIIDSAR SAYQEAMDISKKEMPPTNPIRLGLALNFSVFHYEIANSPEEAISLAKTTFDEAMADLHTLSEDSYKDSTLIMQLLR

DNLTLWTADNAGEEGGEAPQEPQS

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \square \hat{A} \mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than $0.1 \text{ ng/}\tilde{A} \parallel \hat{A} \mu g$ (1 IEU/ $\tilde{A} \parallel \hat{A} \mu g$) as determined by LAL test.