

## 32-7155: Recombinant Human Microtubule-Associated Protein Tau/MAPT-D/Tau-D (C-6His)

**Gene :** MAPT  
**Gene ID :** 4137  
**Uniprot ID :** P10636

### Description

Source: E.coli.  
MW :15.37kD.

Recombinant Human Microtubule-Associated Protein Tau-D is produced by our E.coli expression system and the target gene encoding Gln249-Gln381 is expressed with a 6His tag at the C-terminus. Microtubule-Associated Protein TAU is abundantly expressed in neurons of the central nervous system and less commonly expressed elsewhere, but is also expressed at very low levels in CNS astrocytes and oligodendrocytes. Tau interacts with tubulin to stabilize microtubules and promotes tubulin assembly into microtubules. The C-terminus of TAU binds axonal microtubules while the N-terminus binds neural plasma membrane components, suggesting that tau acts as a linker protein. When tau is defective, and no longer stabilize microtubules properly, it can result in dementias such as Alzheimer's disease and other tauopathies.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** MQIVYKPVDSLKVTSCGSLGNIHHKPGGGQVEVKSEKLDKDRVQSKIGSLDNITHVPGGGNKKIETHKLTFR ENAKAKTDHGAEIVYKSPVVGSDTSPRHLSNVSTGSIDMVDPQLATLADEVSAKQLEHHHHHH

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.