

32-7268: Recombinant Human Histone Chaperone ASF1A/ASF1A (C-6His, N-T7 tag)

Gene : ASF1A
Gene ID : 25842
Uniprot ID : Q9Y294

Description

Source: E.coli.
MW :25.4kD.

Recombinant Human Histone Chaperone ASF1A is produced by our E.coli expression system and the target gene encoding Met1-Met204 is expressed with a T7 tag at the N-terminus, 6His tag at the C-terminus. Human Histone Chaperone ASF1A (ASF1A) belongs to the H3/H4 family of histone chaperone proteins. ASF1A is ubiquitously expressed in many cells and tissues, interacting with histones H3 and H4. ASF1A cooperates with Chromatin Assembly Factor 1 to promote replication-dependent chromatin assembly and with HIRA to promote replication-independent chromatin assembly. In addition, ASF1A is necessary for the formation of senescence-associated heterochromatin foci (SAHF) and efficient senescence-associated cell cycle exit.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM TrisHCl, 1mM DTT, 150mM NaCl, pH 8.0.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : MASMTGGQQMGRGSMKVVQVNNVVLDNPSFPYNPFFQFEITFECIEDLSEDLWKIIVGSAESE EYDQVLDSVLVGPVPAGRHMVFVQADAPNPGLIPDADAVGVTVVLTCTYRGQEFIRVGYYVNN EY TETELRENPPVKPDFSKLQRNILASNPRVTRFHINWEDNTEKLEDAESSNPNLQSLSTDALPSASK GWSTSENSLVNMLESHMDCMLEHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.