

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-7274: Recombinant Human Inhibin beta C Chain/INHBC (N-6His)

Gene ID: 3626 Uniprot ID: P55103

## **Description**

Source: E.coli. MW:14.83kD.

Recombinant Human Inhibin beta C Chain is produced by our E.coli expression system and the target gene encoding Gly237-Ser352 is expressed with a 6His tag at the N-terminus. Inhibins/activins are involved in regulating a number of diverse functions such as hypothalamic and pituitary hormone secretion, gonadal hormone secretion, germ cell development and maturation, erythroid differentiation, insulin secretion, nerve cell survival, embryonic axial development or bone growth, depending on their subunit composition. Inhibins appear to oppose the functions of activins, Inhibins and activins inhibit and activate, respectively, the secretion of follitropin by the pituitary gland.

## **Product Info**

**Amount :** 10 μg / 50 μg

**Content:** Lyophilized from a 0.2 µm filtered solution of 4mM HCl, 1mM DTT.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

**Storage condition :** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: MGSSHHHHHHSSGLVPRGSHMGIDCQGGSRMCCRQEFFVDFREIGWHDWIIQPEGYAMNFCIGQCPLHIAG

MPGIAASFHTAVLNLLKANTAAGTTGGGSCCVPTARRPLSLLYYDRDSNIVKTDIPDMVVEACGCS

## **Application Note**

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\tilde{A} \square \hat{A} \mu g/ml$ . Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin**: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$  (1 IEU/ $\tilde{A} \square \hat{A} \mu g$ ) as determined by LAL test.